


AQA AS CHEMISTRY

# AMOUNT OF SUBSTANCE




1 A sample of 2.18 g of oxygen gas has a volume of 1870 cm<sup>3</sup> at a pressure of 101 kPa.

What is the temperature of the gas?  
The gas constant is  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ .

- A 167 K
- B 334 K
- C 668 K
- D 334 000 K

(Total 1 mark)

2 An ester is hydrolysed as shown by the following equation.



What is the percentage yield of RCOOH when 0.50 g of RCOOH ( $M_r = 100$ ) is obtained from 1.0 g of RCOOR' ( $M_r = 150$ )?

- A 33%
- B 50%
- C 67%
- D 75%

(Total 1 mark)

3 A saturated aqueous solution of magnesium hydroxide contains  $1.17 \times 10^{-3}$  g of  $\text{Mg}(\text{OH})_2$  in 100 cm<sup>3</sup> of solution. In this solution, the magnesium hydroxide is fully dissociated into ions.

What is the concentration of  $\text{Mg}^{2+}(\text{aq})$  ions in this solution?

- A  $2.82 \times 10^{-2} \text{ mol dm}^{-3}$
- B  $2.01 \times 10^{-3} \text{ mol dm}^{-3}$
- C  $2.82 \times 10^{-3} \text{ mol dm}^{-3}$
- D  $2.01 \times 10^{-4} \text{ mol dm}^{-3}$

(Total 1 mark)

**4**

Magnesium reacts with hydrochloric acid according to the following equation.



A student calculated the minimum volume of  $2.56 \text{ mol dm}^{-3}$  hydrochloric acid required to react with an excess of magnesium to form  $5.46 \text{ g}$  of magnesium chloride ( $M_r = 95.3$ ).

Which of the following uses the correct standard form and the appropriate number of significant figures to give the correct result of the calculation?

A  $4.476 \times 10^{-2} \text{ dm}^3$

B  $4.48 \times 10^{-2} \text{ dm}^3$


C  $4.50 \times 10^{-2} \text{ dm}^3$

D  $44.8 \times 10^{-3} \text{ dm}^3$

(Total 1 mark)

**5**

In an experiment to identify a Group 2 metal (X),  $0.102 \text{ g}$  of X reacts with an excess of aqueous hydrochloric acid according to the following equation.



The volume of hydrogen gas given off is  $65 \text{ cm}^3$  at  $99 \text{ kPa}$  pressure and  $303 \text{ K}$ .

The gas constant is  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ .

Which is X?

A Barium

B Calcium


C Magnesium

D Strontium

(Total 1 mark)

6

The following equation represents the oxidation of vanadium(IV) ions by manganate(VII) ions in acid solution.



What volume of 0.020 mol dm<sup>-3</sup> KMnO<sub>4</sub> solution is required to oxidise completely a solution containing 0.010 mol of vanadium(IV) ions?

- A 10 cm<sup>3</sup>
- B 25 cm<sup>3</sup>
- C 50 cm<sup>3</sup>
- D 100 cm<sup>3</sup>

(Total 1 mark)

7

A sample of hydrated nickel sulfate (NiSO<sub>4</sub>.xH<sub>2</sub>O) with a mass of 2.287 g was heated to remove all water of crystallisation. The solid remaining had a mass of 1.344 g.

- (a) Calculate the value of the integer *x*.  
Show your working.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(4)

- (b) Suggest how a student doing this experiment could check that all the water had been removed.

.....  
.....  
.....  
.....

(2)  
(Total 6 marks)

8

Some airbags in cars contain sodium azide ( $\text{NaN}_3$ ).

- (a) Sodium azide is made by reacting dinitrogen monoxide gas with sodium amide ( $\text{NaNH}_2$ ) as shown by the equation.




Calculate the mass of sodium amide needed to obtain 550 g of sodium azide, assuming there is a 95.0% yield of sodium azide.  
Give your answer to 3 significant figures.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

(5)

- (b) If a car is involved in a serious collision, the sodium azide decomposes to form sodium and nitrogen as shown in the equation.



The nitrogen produced then inflates the airbag to a volume of  $7.50 \times 10^{-2} \text{ m}^3$  at a pressure of 150 kPa and temperature of 35 °C.

Calculate the minimum mass of sodium azide that must decompose.

(The gas constant  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ )

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

**(6)**

- (c) Sodium azide is toxic. It can be destroyed by reaction with an acidified solution of nitrous acid ( $\text{HNO}_2$ ) as shown in the equation.



- (i) A  $500 \text{ cm}^3$  volume of the nitrous acid solution was used to destroy completely 150 g of the sodium azide.


Calculate the concentration, in  $\text{mol dm}^{-3}$ , of the nitrous acid used.

.....  
.....  
.....  
.....  
.....  
.....

**(3)**

- (ii) Nitrous acid decomposes on heating.

Balance the following equation for this reaction.



**(1)**

- (d) Sodium azide has a high melting point.

Predict the type of bonding in a crystal of sodium azide.

Suggest why its melting point is high.

Type of bonding .....

Reason for high melting point .....

.....  
.....  
.....  
.....

**(3)**

(e) The azide ion has the formula  $\text{N}_3^-$

- (i) The azide ion can be represented as  $\text{N} \equiv \text{N} - \text{N}^-$   
One of these bonds is a co-ordinate bond.

On the following diagram, draw an arrowhead on one of the bonds to represent the direction of donation of the lone pair in the co-ordinate bond.



(1)

- (ii) Give the formula of a molecule that has the same number of electrons as the azide ion.

.....

(1)

- (iii) Which is the correct formula of magnesium azide?

Tick (✓) **one** box.

$\text{Mg}_3\text{N}$

$\text{MgN}$

$\text{MgN}_6$

$\text{Mg}_3\text{N}_2$

(1)  
(Total 21 marks)



**9**

Zinc is similar to Group 2 metals and forms compounds containing Zn<sup>2+</sup> ions.

Write an equation for the thermal decomposition of zinc carbonate to zinc oxide.

Calculate the percentage atom economy for the formation of zinc oxide from zinc carbonate in this reaction.

Equation .....

Percentage atom economy .....

**(Total 3 marks)**

**10**

Calamine lotion can contain a mixture of zinc carbonate and zinc oxide in suspension in water. A manufacturer of calamine lotion claims that a sample contains 15.00 g of zinc carbonate and 5.00 g of zinc oxide made up to 100 cm<sup>3</sup> with distilled water.

- (a) A chemist wanted to check the manufacturer's claim. The chemist took a 20.0 cm<sup>3</sup> sample of the calamine lotion and added it to an excess of sulfuric acid. The volume of carbon dioxide evolved was measured over time. The chemist's results are shown in the table.

<b>Time / s</b>	0	15	30	45	60	75	90	105	120	135
<b>Volume / cm<sup>3</sup></b>	0	135	270	380	470	530	560	570	570	570


- (i) Plot a graph of the results in the table on the grid. The volume should be on the y-axis. Draw a best-fit curve through **all** the points.

**(3)**

- (ii) Estimate the time taken for the reaction to be completed.

.....

**(1)**



- (b) (i) The volume of carbon dioxide in part (a) was measured at 293 K and at a pressure of 100 kPa.

Use information from your graph to calculate the maximum amount, in moles, of carbon dioxide evolved from the zinc carbonate in this 20.0 cm<sup>3</sup> sample.

The gas constant,  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

Show your working.

.....  
.....  
.....  
.....  
.....  
.....

(3)

- (ii) Use your answer to part (i) to calculate the mass of zinc carbonate in the 20.0 cm<sup>3</sup> sample of calamine lotion.

(If you were unable to complete part (i), you may assume that the amount of carbon dioxide evolved was 0.0225 mol. This is **not** the correct answer.)

.....  
.....  
.....  
.....  
.....

(2)

- (iii) Calculate the difference between your answer to part (ii) and the manufacturer's claim that there are 15.00 g of zinc carbonate in 100 cm<sup>3</sup> of the calamine lotion.

Express this difference as a percentage of the manufacturer's claim.

(If you were unable to complete part (ii), you may assume that the mass of zinc carbonate in the 20 cm<sup>3</sup> sample of calamine lotion was 2.87 g. This is **not** the correct answer.)

Difference .....

Percentage .....

.....

(2)

- (c) Draw a diagram of a suitable apparatus needed to perform the experiment outlined in part (a). Include in your diagram a method for collecting and measuring the carbon dioxide. The apparatus should be airtight.

**(2)**

**(Total 13 marks)**